

**The Incidence and Risk Factors Associated with Stroke Among Patients Who Attended the Medical Ward of Mettu Karl Referral Hospital, South West Oromia, Ethiopia: A Prospective Observational Study**

**Gudisa Bereda<sup>1</sup>, Gemechis Beeda<sup>2</sup>**

doi: <https://doi.org/10.37745/ijeaii.15/vol8n12031>

Published November 23 2023

**Citation:** Bereda G. and Beeda G. (2023) The Incidence and Risk Factors Associated with Stroke Among Patients Who Attended the Medical Ward of Mettu Karl Referral Hospital, South West Oromia, Ethiopia: A Prospective Observational Study, *International Journal of Ebola, AIDS, HIV and Infectious Diseases and Immunity*, Vol.8, No.1, pp.20-31

**ABSTRACTS:** *Stroke as a clinical syndrome typified by “rapidly developing clinical signs of focal or global disturbance of cerebral function, lasting more than 24hours or leading to death, with no apparent cause apart that of vascular origin. Stroke is a major cause of death and disability in many countries. If left untreated, stroke is a debilitating disease that can lead to death. To ascertain the incidence and risk factors associated with stroke among patients who attended the medical ward of Mettu Karl Referral Hospital. A prospective observational study design was carried out from March 13 /2021 to May 19/2021.Data was collected through employing well-structured questioner, and then the collected data was cleared, coded and analyzed by Statistical Packages for Social Sciences 25.0 version statistical software. The relationship between risk factors and stroke subtypes were examined by logistic regression and chi -square test. Statistical significant was P value at <0.05. From the 84 patients, nearly above one-half 47(56.0%) of participants were male patients. Nearly one half 39(46.4%) of respondents age were between 45-65 years. From the study subjects, majority 48(57.1%) of respondents were have ischemic type of stroke followed by 29(34.5%) haemorrhagic stroke. In multivariable regression analysis, the age>65 years( $\chi^2=3.58,p=0.001$ ),male( $\chi^2=1.78,p0.09$ ),comorbidities( $\chi^2=2.15,p=0.005$ ),medicationadherence( $\chi^2=4.67,p=0.001$ ),don't reduce salt in diet ( $\chi^2=,p=0.076$ ), eating fatty diet( $\chi^2=1.79,p=0.005$ ), and rural residents( $\chi^2=2.05, p=0.006$ ) were significantly associated with stroke risk factors. Among the study participants, less than half of patients were smoke cigarette every, were smoking cigarette after diagnosed, were do regularly exercise regularly, were drink alcohol everywhere drink alcohol currently, were don't reduce salt in diet, and were eat fatty diet. Health care workers should have to treat stroke risk factors to manage the stroke mortality rate.*

**KEYWORDS:** Stroke, Risk factors, Medical ward, Mettu Karl Referral Hospital, Ethiopia.

## INTRODUCTION

Stroke is defined as a rapidly developing global or focal neurological deficit maintaining more than 24 h or causing death without clear causes other than vascular

Publication of the European Centre for Research Training and Development -UK

origin". Also, stroke can be defined as a rapid damage in brain function caused by brain arteries decrease or stop [1]. It is divided into three types; most commonly hemorrhagic, ischemic, and Transient Ischemic Attacks (TIA). Stroke is a major public health problem, affecting millions of people in both developed and developing countries. It mainly occurs because of poor blood flow to the brain which results in cell death. Inability to move, cognitive impairment, talking problems, feeling as the world is spinning, severe headache and loss of vision are key signs and symptoms of stroke [2]. The incidence, prevalence, and mortality rate of stroke have been increased worldwide, with most of the burden being in low and middle-income countries. The incidence, prevalence, and mortality rate of stroke have been increased worldwide, with most of the burden being in low and middle-income countries [3]. Developing countries have a higher burden of non-communicable diseases than the rest of the world. More than two-third (70%) of strokes occur in low- and middle-income countries [4]. Globally, stroke has covered about 16 million incidences and 9.7% deaths. Over 23 and 7.8 million new stroke cases and deaths are expected by 2030, respectively if there will not be intervention [5]. According to Global Burden of Disease (GBD) study in 2010, more than 11 million ischemic strokes occurred while 63% of them were in lowland middle-income countries. Also, near 3 million deaths occurred due to ischemic stroke. About 13% of stroke is of a hemorrhagic type [6]. According to GBD study in 2010, there were about 5.3 million hemorrhagic stroke cases, out of which about 80% occurred in low- and middle-income countries. Over 3 million deaths occurred from hemorrhagic stroke [7]. Stroke in sub-Saharan Africa is a major public health problem, with higher mortality than in developed countries and occurrence at a younger age [8]. The American heart association plans to reduce ease and deaths from stroke by 20 percent in 2020 by focusing on seven key health factors and behaviors that decrease the risk of stroke, those are not-smoking, physical activity, healthy diet, body weight, and control of cholesterol, blood pressure, and blood sugar [9]. Stroke has different risk factors, which can be grouped into modifiable and nonmodifiable risk factors. Major risk factors for stroke include age, history of cerebrovascular event, smoking, alcohol consumption, physical inactivity, hypertension, dyslipidemia, diabetes mellitus, cardiovascular diseases, obesity, metabolic syndrome, diet, nutrition, and genetic risk factors [10]. The most factors linked to intracerebral hemorrhage are HTN and oral anticoagulation therapy; while, smoking, alcohol consumption, and HTN have been reported as risk factors to subarachnoid hemorrhage [11].

According to the World Health Organization, the second cause of death over the past 2002-2012 has been stroke. Also, it has estimated that in 2004 about 5.7 million deaths happened because of stroke which accounted for 9.7% of deaths; more than 85% of these deaths occurred in middle-income and low income countries [12]. The reasons for this trend are not clear, but it can be attributed to getting more information about the risk factors of stroke. There were several studies which confirm this relationship. One of them conducted in Saudi Arabia in (1993) reported hypertension as one of the most serious stroke risk factors between the Saudi populations. Thus far, no nationwide has been conducted recently on the prevalence and incidence of strokes in Saudi Arabia.

However, it was reported in a study that the rate of stroke in Saudi Arabia was 29.8/100,000. They also reported that ischemic strokes (69%) predominated and Subarachnoid Hemorrhage (SAH) accounting only for 1.4% [13]. Developing countries have shared the majority of global stroke mortality [14]. About 85.5% of world's stroke deaths have occurred in developing countries. There was an increment of stroke prevalence by 8% in Africa between 2002 and 2004. Smoking, alcohol intake, diabetes mellitus, poor physical activity and unhealthy diets were the stated factors to stroke. Old age also has a positive linkage with stroke [15]. In 2015/16-2019/20 the Ethiopian health sector development program projects to decrease by 12.5% premature mortality from non-communicable disease (NCDs). In Ethiopia currently, stroke is one of the greatest public health problems, accounts for 7% of total deaths. Study in Mekelle, Ethiopia showed that the stroke was the third most common cause of medical intensive care unit admission (15.2%) and the first cause of death, which accounts for 17% of all deaths in the medical intensive care unit [16]. Similarly, hypertension is responsible for 66.2% of all stroke admission and 38% of all stroke were on anti-hypertensive treatment [17]. This study was conducted to estimate the magnitude of risk factors of stroke and mortality rate in hospital and, the unavailability of a nationwide study that determines the magnitude of risk factors and mortality rate in-hospital of stroke is an important research gap.

## METHODS

**Study setting:** The study was conducted in MKRH, Mettu town, South West Oromia, Ethiopia which is found at 600 km from Addis Ababa. There are different wards and clinics within MKRH; those include internal medicine ward, surgery ward, pediatric ward, gynecology and obstetrics ward, Ante natal clinic, dental clinics, tuberculosis clinic, anti-retroviral therapy clinic and ophthalmologic clinic.

**Study design and period:** A Hospital based prospective observational study design was conducted from March 13 /2021 to May 19/2021.

**Inclusion and exclusion criteria:** Participants, aged 18 over (patients, patients' companions and visitors), attending urban primary health care centers held as part of the study to benefit from preventive or curative care, were included in the study. Health workers, Pregnant mothers, Patients who admitted to intensive care unit were excluded.

**Sample size determination and recruitment of study participants:** The sample size was determined by using the Single Population proportion Formula: The sample size was determined based on "P" value which was taken from Ethiopia, P=0.151, or 15.1%.  
$$n = \frac{(Z_{\alpha/2})^2 \cdot P(1-P)}{d^2}$$
  
n= sample size, P=prevalence of stroke, d=margin of sampling error tolerated, z=the standard normal value at confidence interval of 95%.  $n = \frac{(1.96)^2(1-0.151) \times (0.151)}{(0.05)^2} = 197$ . Since the total number of stroke patients was less than 10,000, reduction formula (correction formula) was applied as follow;  $n_f = n / (1 + (n/N))$ ,

Publication of the European Centre for Research Training and Development -UK

$n_f = 197 / (1 + (197/124)) = 76$ . When 10% contingency is added to minimize non response rate. then final sample size was found to be 84. Purposive sampling technique was used to recruit samples for the study in each day of the data collection process until the desired sample size was obtained.

**Data collection tools and procedures:** A face-to-face questionnaire survey was used for data collection of respondents with the first part including sections reserved for: socio-demographic characteristics (age, sex, marital status, level of education, spoken languages, place of residence, socioeconomic level (family income), Behavioral factors (smoking status, exercise per day, and dietitians (which restricted their intakes of sugar, salt, and fat), Clinical characteristics (family histories of strokes, the duration of disease, and self-reported comorbidities), Medications history and associated comorbidities (high blood pressure, diabetes, hypercholesterolemia, cardiac disease, history of stroke in respondent or immediate family, and history of stroke in relatives, acquaintances, or neighbors). During data collection, data collector measures weight, height, blood pressure, as follows. Weight was measured in light clothing and without shoes by calibrated UNICEF Seca digital weighing scale. Stadiometer in centimeter in erect position at a precision of 0.5 cm without shoes was used to measure height. Mercury sphygmomanometer was used to measure blood pressure average of two measurements 5 minutes apart was recorded for those who we take BP during the data collection.

**Data quality assurance:** In order to assure quality of data important measures were undertaken including: the patient card number was used, to check for if there is invalid and incomplete pertinent response and these cards were also coded so over or under count was not matter. The data collected was checked for completeness and consistency on daily basis.

**Data management and statistical analysis:** The gathered quantitative data was cleared, arranged, coded, and then analyzed through employing SPSS version 25.0 statistical Software. Categorical variables were expressed by percentage and frequency, whereas continuous variables were present by mean and standard deviation. For comparison of baseline variables between groups, chi-square tests were performed for categorical data, and Student's *t*-tests was carried out for continuous data. Bivariate and multivariate logistic regression analysis was computed to identify factors associated with the stroke. All independent variables with a p-value <0.25 in the uni-variate analysis were taken into account in the multivariate logistic regression analysis. P values <0.05 was considered to indicate statistical significance.

**Ethical Clearance:** Ethical clearance was taken from Negelle Borena, college of health sciences, and informed consent was obtained from each subject. No any personal identifier was taken from the patient registration log book. The information was recorded anonymously and behind the scenes and beneficence were assured throughout the study period.

**Operational Definitions**

**Stroke** is the abrupt onset of a neurologic deficit and attributable to a focal vascular cause.

**Ischemic stroke:** is blockage artery that supplies oxygen rich blood to the brain.

**Hemorrhagic stroke:** is leakage of artery in the brain.

**RESULTS**

Socio-demographic characteristics of the study participants.

From the 84 patients, nearly above one-half 47(56.0%) were male patients. Nearly one half 39(46.4%) of respondents age were between 45-65 years, and majority 55 (65.5%) of patients were earn monthly income <1000 ETB. More than half 51(60.7%) of participants were married, 43(51.2%) were uneducated, and 49(58.3%) were dwell in rural area. Less than half 38(45.2%) of participants had at least one social habit.

Table 1. Socio-demographic characteristics of the stroke patients at MKRH, South western, Ethiopia. 2021 (n=84)

Variables	Category	Frequency	Percent
Age	<45 years	24	28.6
	45-65 years	39	46.4
	>65 years	21	25.0
Sex	Male	47	56.0
	Female	37	44.0
Monthly income	<1000 ETB	55	65.5
	≥1000 ETB	29	34.5
Marital status	Married	51	60.7
	Unmarried	33	39.3
Educational status	Educated	41	48.8
	Uneducated	43	51.2
Residency	Urban	49	58.3
	Rural	35	41.7
Social habit	Yes	38	45.2
	No	46	54.8

**Behavioral factors of the respondents**

Among the study participants, less than half 30(35.7%) of patients were smoke cigarette every, 36(42.9%) were smoking cigarette after diagnosed, 34(40.5%) were do regularly exercise regularly, 29(34.5%) were drink alcohol every, 25(29.8%) were drink alcohol current, 23(27.4%) were reduce salt in diet, and 34(40.5%) were eat fatty diet. The medication non-adherence was 32(38.1%).

Table 2. Behavioral factors of the stroke patients at MKRH, South western, Ethiopia. 2021 (n=84)

Variables	Category	Frequency	Percent
Have you ever smoke cigarette	Yes	30	35.7
	No	54	64.3
Smoking after you diagnosed	Yes	36	42.9
	No	48	57.1
Regular physical exercise	Yes	34	40.5
	No	50	59.5
Medication non-adherence	Yes	32	38.1
	No	52	61.9
Have you ever drink alcohol	Yes	29	34.5
	No	55	65.5
Current alcohol drink	Yes	25	29.8
	No	59	70.2
Do you reduce salt in diet	Yes	23	27.4
	No	61	72.6
Do you eat fatty diet	Yes	34	40.5
	No	50	59.5

## Clinical and drug related characteristics of the patients

In our survey, slightly less than half 41(48.8%) of the study participants were have laboratory values, and more than half 52(61.9%) of the respondents had at least one comorbidities. A majority 27(32.1%) of participants BMI were between 18.5-24.99kg/m<sup>2</sup>. Above half 45(53.6%) of the participants were have past medical history and half 42(50.0%) of study subjects were have past medication history. Less than half 37(44.0%) of participants had past family medical history, and 35(41.7%) participants had past family medication history. From the study subjects, majority 48(57.1%) were have ischemic type of stroke followed by 29(34.5%) haemorrhagic stroke. Majority 50(59.5%) of study subjects had high level blood pressure and only 32(38.1%) of patients had family history of stroke.

Table 3. Clinical and drug related characteristics of the stroke patients at MKRH, South western, Ethiopia. 2021 (n=84)

Variables	Category	Frequency	Percent
Laboratory values	Yes	41	48.8
	No	43	51.2
Comorbidities	Yes	52	61.9
	No	32	38.1
BMI	<18.5kg/m <sup>2</sup>	20	23.8
	18.5-24.99kg/m <sup>2</sup>	27	32.1
	25-29.9kg/m <sup>2</sup>	18	21.4

## Publication of the European Centre for Research Training and Development -UK

	$\geq 30\text{kg/m}^2$	19	22.6
Past medical history	Yes	45	53.6
	No	39	46.4
Past medication history	Yes	42	50.0
	No	42	50.0
Past family medical history	Yes	37	44.0
	No	47	55.0
Past family medication history	Yes	35	41.7
	No	49	58.3
Type of stroke	Haemorrhagic	29	34.5
	Ischemic	48	57.1
	Transit	5	6.0
	Unclassified	2	2.4
Blood pressure	Normal	34	40.5
	High level	50	59.5
Family history of stroke	Yes	32	38.1
	No	452	61.9

## Comorbidities of the stroke among stroke patients

From the total of 84 study subjects 14(16.7%) of patients were have hypertension followed by acute kidney injury 12(14.35%) and chronic kidney disease 10(11.9%). Anemia 6(7.1%) were least identified comorbidities followed by 5(6.0%) seizure and 4(4.8%) asthma.

Table 4. Comorbidities of the stroke patients at MKRH, South western, Ethiopia. 2021 (n=84)

Variables	Category	Frequency	Percent
Comorbidities	Hypertension	14	16.7
	Ischemic heart disease	7	8.3
	Diabetes mellitus	9	10.7
	Chronic kidney disease	10	11.9
	Acute kidney injury	12	14.3
	Seizure	5	6.0
	Anemia	6	7.1
	Asthma	4	4.8
	Dyslipidemia	8	9.5
	Stroke	7	8.3
	Others	2	2.4

Logistic regression analysis for factors associated with stroke risk factors

Publication of the European Centre for Research Training and Development -UK

In multivariable regression analysis, the, >65 years, male, high BMI, comorbidities, medication non-adherence, do you reduce salt in diet, and do you eat fatty diet were significantly associated with stroke risk factors. Participants were age above >65 years were 3.57 times more likely stroke risk factors(AOR:3.579;95%CI:2.931-7.208; P=0.001),and male were 1.78 times more likely stroke risk factors (AOR: 1.78; 95%CI:1.935-2.916;P=0.09).Comorbidities were 2.15 times more likely stroke risk factors(AOR:2.15; 95%CI:2.074-3.858;P=0.005) and patients who have medication non-adherence were 2.15 times more likely stroke risk factors (AOR:4.67;95%CI: 4.524-9.105;P=0.001).Don't reduce salt in diet were 1.79 times more likely stroke risk factors (AOR:1.79;95%CI:1.971-2.431;P=0.076) and eating fatty diet were 2.05 times more likely stroke risk factors (AOR:1.79;95%CI: 2.157- 4.381;P=0.005).

Table 5. Factors associated with stroke patients at MKRH, South western, Ethiopia. 2021 (n=84)

Variables	Category	n(%)	AOR (95% C.I)	P-value
Age	<45 years	24(28.6)	1	
	45-65 years	39(46.4)	1.32(1.045-1.453)	0.734
	>65 years	21(25.0)	3.579(2.931-7.208)	0.001
Sex	Female	37(44.0)	1	
	Male	47(56.0)	1.78(1.935-2.916)	0.09
Blood pressure	Normal	34(40.5)	1	
	High level	50(59.5)	0.97(0.057-1.032)	0.317
Comorbidities	Yes	52(61.9)	1	
	No	32(38.1)	2.15(2.074-3.858)	0.005
Medication non-adherence	Yes	32(38.1)	1	
	No	52(61.9)	4.67(4.524-9.105)	0.001
Have you ever smoke cigarette	Yes	30(35.7)	1	
	No	54(64.3)	1.16(1.128-1.976)	0.027
Have you ever drink alcohol	Yes	29(34.5)	1	
	No	55(65.5)	0.92(0.023-1.032)	0.87
Do you reduce salt in diet	Yes	23(27.4)	1	
	No	61(72.6)	1.79(1.971-2.431)	0.076
Do you eat fatty diet	No	50(59.5)	1	
	Yes	34(40.5)	2.05(2.157-4.381)	0.005



## **DISCUSSION**

According to WHO data published in 2017, stroke deaths in Ethiopia reached 6.23% of total deaths. In addition, the age-adjusted death rate of stroke in the country is 89.82 per 100,000 of the population. Previous reports that showed the future trend of stroke in SSA revealed that stroke burden will increase over the coming years owing to poor healthcare seeking behavior, and poor neurologic interventions [18].

The present study revealed rural residents were 3.74 times more likely higher stroke risk factors (AOR:3.74;95%CI:2.904-7.516;P=0.006) than urban area residents were higher than the study conducted in Iran[19] those patients who were resident in urban districts were more than 1.5 times at higher risk of developing stroke compared with residents of rural areas, which can be probably attributed to the unhealthy lifestyle practices with regard to lack of physical activity and unhealthy diet. Because rural residents report less leisure time physical activity and lower seatbelt use than urban residents and rural residents also have higher rate of poverty, less access to health care and are less likely to have health insurance.

Our survey showed that 14(16.7%) of stroke patients were have hypertension followed by acute kidney injury 12(14.35%) and chronic kidney disease 10(11.9%) comorbidities were in line with the meta-analysis study done in Ethiopia[20] showed diabetes is a well-established risk factor of stroke, and our analysis showed that diabetes mellitus is the second most common comorbidity of stroke. Nearly one-tenth (8%) of stroke patients had diabetes mellitus. Due to hypertension were the risk factors for stroke due to in hypertension blood pressure was high which factors of coronary heart disease and stroke.

The current study showed that nearly above one-half 47(56.0%) were male patients were consistent with the study surveyed in Felege Hiwot Referral hospital[21] which showed male patients were more affected by stroke; 20 (62.5%) in this study. Due to male had no cardio-protective hormone which protect hypertensive heart diseases and men diagnosed with diabetes more than women, which increases the risk of stroke because it can cause disease of blood vessels in brain.

The present survey showed that majority 48(57.1%) of respondents were have ischemic type of stroke followed by 29(34.5%) haemorrhagic stroke were in line with the study conducted in Felege Hiwot Hospital[21] showed that ischemic stroke was more prevalent (56.7%) than haemorrhagic. Because they happen when blood clot blocks the flow of blood and oxygen to the brain and the occlusions contribute to around 85% of casualties of stroke patients.

Our study revealed that participants were age above >65 years were 3.57 times more likely stroke risk factors (AOR:3.579;95%CI:2.931-7.208;P=0.001) were in line with the study conducted in Felege Hiwot Hospital[21] showed that older age (>50 years

Publication of the European Centre for Research Training and Development -UK

old) was factor associated with stroke;AOR=0.41.Due to stroke was a disease of aging and the risk increase with age,the incidence doubling with each decade after the age of 45 years and over 70% of all strokes occur above the age of 65.

The current study revealed that patients who have medication non- adherence were 2.15 times more likely stroke risk factors (AOR:4.67;95%CI:4.524-9.105;P=0.001) were in line with the study conducted in Ayder Comprehensive Specialized Hospital [22]showed that medication none-adhered were 4 times(AOR=3.967,95%CI:2.256–6.973) more likely to be at high risk for stroke than the medication adherent hypertensive patients.Here non adherence to cardiovascular medication perhaps increase of stroke occurrence.

Our study revealed that the patients don't reduce salt in diet were 1.79 times more likely stroke risk factors (AOR:1.79;95%CI:1.971-2.431;P=0.076) and patients eat fatty diet were 2.05 times more likely stroke risk factors (AOR:1.79;95%CI:2.157-4.381;P=0.005) were in line with the study conducted in Ayder Comprehensive Specialized Hospital[22] showed that hypertensive patients who did not reduce salt in diet were around 3.2 times more likely to be at high risk for stroke (AOR=3.249,95% CI:1.544–6.837).Because consuming too much salt can increase the risk of having high blood pressure ,which is a risk factors for coronary heart disease and stroke and eating unhealthy (fat) diet can increase your chances of having a stroke because it may lead to an increase in the blood pressure and cholesterol levels.

Our study revealed that male stroke patients were 1.78 times more likely stroke risk factors (AOR:1.78;95%CI:1.935-2.916;P=0.09) than females and patients who have comorbidities were 2.15 times more likely stroke risk factors (AOR:2.15;95%CI: 2.074-3.858;P=0.005) than with out comorbidities.

## **CONCLUSION**

Among the study participants, less than half of patients were smoke cigarette every, were smoking cigarette after diagnosed, were do regularly exercise regularly, were drink alcohol every, were drink alcohol current, were reduce salt in diet, and were eat fatty diet. The medication non-adherence of stroke patients was high. Patients age >65 years, male, comorbidities, medication non- adherence, don't reduce salt in diet, and eating fatty diet were significantly associated with stroke risk factors.

## **Acknowledgments**

We had extended our grateful thank to the study participants

## **REFERENCES**

- 1.Hof P,Mobbs C.Handbook of the neuroscience of aging.London,Academic Press,2009:p.485

Publication of the European Centre for Research Training and Development -UK

2. Longo DL, Kasper DL, Jameson JL, Fauci AS, et al. Harrison's Principles of Internal Medicine 18th Edition, Volume 1. <http://harrell.library.psu.edu/c.php?>
3. Yan LL, Li C, Chen J, Miranda JJ, et al. Prevention, management, and rehabilitation of stroke in low- and middle-income countries. *Neurological Sci.* 2016;2:21–30
4. World Stroke Organization. Facts and Figures about Stroke. <https://www.world-stroke.org/component/content/article/16-forpatients/84-facts-andfigures-about-stroke>. Accessed 18 Nov 2019
5. Howard VJ. Reference module in neuroscience and biobehavioral psychology; 2018.
6. D. Mozafarian, E. J. Benjamin, A. S. Go, D. K. Arnett, et al., "Heart disease and stroke statistics—2015 update: a report from the American Heart Association," *Circulation*, vol. 131, no. 4, pp. e29–e322, 2015.
7. Krishnamurthi, A. E. Moran, M. H. Forouzanfar et al., "Teglobal burden of hemorrhagic stroke: a summary of findings from the GBD 2010 study," *Global Heart*, vol. 9, no. 1, pp. 101–106, 2014.
8. Kalapo K O, Vento S. Stroke: a realistic approach to a growing problem in sub-Saharan Africa is urgently needed. *Tropical Medicine & International Health* 2011;16: 707–10.
9. Benjamin E. J., et al., "Heart disease and stroke statistics—2017 update: a report from the American Heart Association." *Circulation*, 2017. 135(10): p. e146–e603. <https://doi.org/10.1161/CIR.0000000000000485> PMID: 28122885
10. Hopewell J. C. and R. Clarke, "Emerging Risk Factors for Stroke: What Have We Learned from Mendelian Randomization Studies?" *Stroke*, vol. 47, no. 6, pp. 1673–1678, 2016.
11. von Sarnowski B, J. Putaala, U. Grittner et al., "Lifestyle risk factors for ischemic stroke and transient ischemic attack in young adults in the stroke in young Fabry patients study," *Stroke*, vol. 44, no. 1, pp. 119–125, 2013
12. World Health Organization (WHO). The top 10 causes of death [Internet]. World Health Organization. 2016 [cited 21 April 2016]. Available
13. World Health Organization (WHO). WHO | Risk factors [Internet]. Who.int. 2016 [cited 18 April 2016]. Available from: [http://www.who.int/topics/risk\\_factors/en/](http://www.who.int/topics/risk_factors/en/)
14. Thun MJ, Carter BD, Feskanich D, et al. 50-Year Trends in Smoking-Related Mortality in the United States. *N Engl J Med.* 2013;368:4,351–64.
15. Seshadri S, Beiser A, Kelly-Hayes M, Kase CS, Au R, Kannel WB, Wolf PA. The Lifetime Risk of Stroke: Estimates from the Framingham Study. *Stroke.* 2006; 37:2,345–35
16. Chhabra M, Gudi SK, Rashid M, et al. Assessment of knowledge on risk factors, warning signs, and early treatment approaches of stroke among community adults in North India: A telephone interview survey. *Journal of Neurosciences in Rural Practice.* 2019;10(3):417–422. DOI: <https://doi.org/10.1055/s-0039-1697561>
17. Gebremariam S. A. and Yang H. S., Types, risk profiles, and outcomes of stroke patients in a tertiary teaching hospital in northern Ethiopia. *eNeurological Sci.* 2016.3: p. 41–47. <https://doi.org/10.1016/j.ensci.2016.02.010> PMID: 29430535.

Publication of the European Centre for Research Training and Development -UK

18. BeLue R, Okoror TA, Iwelunmor J, Taylor KD et al. An overview of cardiovascular risk factor burden in sub-Saharan African countries: a socio-cultural perspective. *Glob Health*. 2009;5(1):10
19. K. Teo, S. Lear, S. Islam et al., "Prevalence of a healthy lifestyle among individuals with cardiovascular disease in high-, middle- and low-income countries: The Prospective Urban Rural Epidemiology (PURE) study," *Journal of the American Medical Association*, vol. 309, no. 15, pp. 1613–1621, 2013.
20. Alene M, Agazhe M, Assemie Y, Yismaw L, and Bekele D Ketema. Magnitude of risk factors and in-hospital mortality of stroke in Ethiopia: a systematic review and meta-analysis. *Alene et al. BMC Neurology* (2020) 20:309 <https://doi.org/10.1186/s12883-020-01870-6>
21. Mulat B, Mohammed J, Yeseni M, Alamirew M. Magnitude of stroke and associated factors among patients who attended the medical ward of Felege Hiwot Referral Hospital, Bahir Dar town, Northwest Ethiopia. *Ethiop. J. Health Dev.* 2016
22. Hagos H, Mekonen M, Molla M, Birhanu B, Belete T, Mossie T, Tsegabrhan H, Gebreslassie. Factors associated with stroke among adult patients with hypertension in Ayder Comprehensive-Specialized Hospital, Tigray, Ethiopia, 2018: A case-control study. February 13, 2020. | <https://doi.org/10.1371/journal.pone.0228650>